

Norac Concepts Inc.
P.O. Box 31097
Guelph, ON N1H 8K1
519-821-3633

GROUP	M	FUNGICIDE
-------	---	-----------

AGRICULTURAL
Net Content: 10 kg

TUBERSEAL
Potato Seed Piece Dust Contains Douglas Fir Bark

To Control Fusarium Seed Piece Decay

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

GUARANTEE
mancozeb..... 16%

REGISTRATION NO. 17042 PEST CONTROL PRODUCTS ACT

PRECAUTIONS

Store product in a cool, dry, ventilated place. Do not allow product to become wet or overheated during storage as chemical changes may occur which impair fungicidal effectiveness and may also generate flammable vapours. May irritate eyes, nose, throat and skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating, drinking or smoking. When treating or handling treated seed, work in a well-ventilated area, and wear a suitable dust mask, goggles and gloves. Do not contaminate food or feed. Do not store near food or feed. Do not contaminate any body of water. Treated seed should be labelled "POISONOUS TO HUMANS AND ANIMALS. This seed has been treated with mancozeb for the control of fusarium decay. Do not use for food or feed purposes". If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact 1-866-375-4648 or www.cropro.org.

FIRST AID

If swallowed, induce vomiting and get medical attention. **If on skin**, wash with soap and water. **If in eyes**, flush with water for 5 to 10 minutes and get medical attention. Take container, label, product name and Pest Control Product Registration Number with you when seeking medical attention.

EMERGENCY TELEPHONE NUMBER

All hours, (613) 787-5620 **ONLY** for health and environmental information.

DISPOSAL

Empty container thoroughly into application equipment. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER

This control product is to be used in accordance with the directions on this label. It is an offense under the *Pest Control Products Act* to use a control product under unsafe conditions.

DIRECTIONS

Apply at the rate of 500 g per 100 kg of seed. Thoroughly coat the surface of whole or cut seed pieces with dust. If treated whole seed is cut, make a second application to protect the cut surfaces. Plant as soon as possible after treating. However, if planting of cut seed is delayed beyond 2 days after treating, seed should be stored in open crates to allow air drying until dry before bagging or loose piling.

This product contains no colourant. An appropriate colourant must be added when this product is applied to seed. Regulations pertaining to the "Seed Act" must be strictly adhered to when using this product.

RESISTANCE MANAGEMENT:

For resistance management, please note that TUBERSEAL contains a Group M fungicide. Any fungal population may contain individuals naturally resistant to TUBERSEAL and other Group M fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of TUBERSEAL or other Group M fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group when such use is permitted.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- If disease continues to progress after treatment with this product, do not increase the use rate.

Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact Norac Concepts at 519-821-3633 or at www.noracconcepts.com.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product, and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.